

1 **MOLECULAR ANALYSIS OF NSP4 CODING GENE OF PORCINE ROTAVIRUS IN**
2 **BRAZIL**
3
4 *(ANÁLISE MOLECULAR DO GENE CODIFICADOR DA NSP4 DE ROTAVÍRUS SUÍNOS*
5 *NO BRASIL)*

6 **SUMMARY**

7

8 The non-structural protein 4 (NSP4) has different roles in rotaviral replication, morphogenesis,
9 and enterotoxin-like activity causing secretory diarrhea. A total of 11 partial nucleotide
10 sequences of NSP4 coding gene were defined from group A rotavirus circulating in Brazilian
11 swine herds. On comparing the viral sequences of diarrheagenic peptide area (amino acid 114-
12 135), there was a single point mutation at amino acid 135 presented by two strains with amino
13 acid alanine, and valine in the others. The NSP4 gene phylogeny showed that all strains
14 clustered into E1 genotype, and the nucleotide identity between Brazilian strains ranged from
15 92.4% and 100%, while the putative amino acid identity, between 95.8% and 100%. As a
16 conclusion, these data demonstrate the occurrence of a common NSP4 genotype described
17 elsewhere in pigs and low diversity between the samples from the surveyed areas.

18

19 **KEY WORDS:** Genotypes. Non-structural protein. Reoviridae.

20

21 **RESUMO**

22

23 A proteína não estrutural 4 (NSP4) desempenha diferentes funções na replicação e na
24 morfogênese dos rotavírus, possuindo, ainda, uma atividade de enterotoxina, causando diarreia
25 do tipo secretória. Um total de 11 sequências parciais de nucleotídeos do gene codificador da
26 NSP4 de rotavírus suínos de criações brasileiras foram definidas como pertencentes ao grupo
27 A. Comparando-se as sequências virais da área do peptídeo toxigênico, que compreende a
28 porção entre os aminoácidos de 114 a 135, constatou-se uma única mutação pontual no
29 aminoácido 135, sendo que duas amostras apresentaram alanina, e as demais, valina. A análise
30 filogenética do gene demonstrou que todas as amostras pertencem ao genotipo E1, e que a
31 identidade nucleotídica das amostras brasileiras variou de 92,4% a 100%, enquanto que a
32 identidade de aminoácidos, de 95,8% a 100%. Assim, esses dados mostram a ocorrência de um
33 genotipo comum da NSP4 já descrito anteriormente em suínos, com uma baixa diversidade
34 entre as amostras encontradas.

35

36 **PALAVRAS-CHAVE:** Genótipos. Proteína não-estrutural. Reoviridae.

37

INTRODUCTION

38

39

40 Group A rotavirus (RV-A), members of the *Reoviridae* family, genus *Rotavirus*, are
41 regarded as a major cause of gastroenteritis both in humans and animals worldwide
42 (KAPIKIAN et al., 2001). The RV-A genome consists of 11 segments of double-stranded RNA,
43 encoding six structural virus proteins (VP1-VP4, VP6 and VP7) and six non-structural proteins
44 (NSP1-NSP6). The complete virus is a triple-layered particle, with VP4 and VP7 constituting
45 the outer layer whose respective encoding genes are markers for P and G genotypes,
46 respectively (KING et al., 2012). So far, 35 genotypes P and 27 G have been defined
47 (MATTHIJNSSENS et al., 2011). The inner capsid protein VP6 important on host immunity
48 and determines groups A-G, and more recently, a novel RV-H has been discovered (KING et
49 al., 2012; MATTHIJNSSENS et al., 2012). Genotypes previously described in pigs include
50 G1, G2, G3, G4, G5, G6, G11 and G12, usually associated with P[6], P[7], P[13], P[19], P[23],
51 P[26], and P[27] (MATTHIJNSSENS et al., 2008b; TONIETTI et al., 2013).

52 The non-structural protein 4 (NSP4), encoded by gene segment 10, has multiple functions
53 in RVs morphogenesis and pathogenesis. It has an enterotoxin-like activity (BALL et al., 1996)
54 and has been identified as a viroporin (HYSER et al., 2012). The peptide 114-135 is considered
55 to trigger a signal transduction pathway as it increases intracellular calcium leading to chloride
56 secretion, and therefore secretory diarrhea, as it has been shown in mice (BALL et al., 1996;
57 HUANG et al., 2004; TIAN et al., 1995). Changes within this region have been associated with
58 alterations in the toxigenic activity of NSP4 and virulence of RVs (BALL et al., 1996; ZHANG
59 et al., 1998). So far 14 NSP4 genotypes have been defined from RV-A samples infecting
60 Human and animal hosts (MATTHIJNSSENS et al., 2008a).

61 The aim of this investigation was to sequence and analyze a partial fragment of NSP4
62 gene of RV-A from different Brazilian pig herds to define their phylogenetic relations with
63 other animal and human isolates described elsewhere.

64

65 MATERIAL AND METHODS

66

67 A total of 11 stool samples from pigs with diarrhea from three cities in São Paulo's State,
68 Brazil, were collected in 2008 and screened with polyacrylamide gel electrophoresis (PAGE),
69 ELISA, and characterized in P and G genotypes as previously described (GOUVEA et al.,
70 1994a,b).

71 Feces suspensions (v/v; 50%) were prepared with phosphate-buffered saline 0.01M, pH
72 7.2, clarified at 5000g/15 min at 4°C, and the supernatants used in the assays.

73 Extraction of total RNA from the reference RVs strain (NCDV) and the supernatants of
74 the field samples were carried out with TRIzol Reagent™ (Invitrogen, Carlsbad, CA, USA)
75 according to the manufacturer's instructions.

76 For RT-PCR, 5.6µL of RNA solution was mixed with 1.4µL of DMSO and denatured at
77 95°C for 5 minutes and kept in ice. Then it was added to a solution of 1x First Strand Buffer
78 (Invitrogen™), 1 mM of each dNTP, 10 mM DTT, and 1 µM of each primer targeting NSP4
79 coding gene (10BEG16 and 10END722) as described by Lee et al. (2000) and 200U of reverse
80 transcriptase (Invitrogen™), to a 13µL final reaction volume. This mixture was then heated at
81 42°C for 1 hour and 70°C for 15 min at thermal cycler.

82 PCR amplification was carried out by adding 5µL of cDNA of the RT reaction in a mix
83 containing 1x PCR Buffer (Invitrogen™), 0.2 mM of each dNTP, 0.5 µM of each primer
84 (10BEG16 and 10END722), as described by Lee et al. (2000), 2 mM of MgCl₂, and 2.5U of
85 Taq DNA Polymerase (Invitrogen™) and ultra-pure water for a final reaction volume of 50 µL.

86 This mixture was heated at 94°C for 2 min, followed by 30 cycles each at 95°C for 45 s, 49°C
87 for 30 s, 72°C for 1.5 min, and one cycle at 72°C for 10 min. The products of the PCR were
88 resolved on a 1.5% agarose gel stained with 0.5µg/mL ethidium bromide.

89 Amplicons of 725 bp in length were purified with Illustra GFX™ PCR DNA and Gel
90 Band Purification Kit, according to the manufacturer's instructions (GE Healthcare) and
91 submitted to bi-directional sequencing with BigDye 3.1™ (Applied Biosystems, Carlsbad, CA,
92 USA) according to the manufacturer's instructions. The reaction products of the sequencing
93 reactions were resolved in the automatic sequencer ABI-377™ (Applied Biosystems, Carlsbad,
94 CA, USA).

95 Nucleotide sequences obtained in this study (nt 66 to nt 566, using as reference Gottfried
96 Strain accession number GU199490) (Table 1) were aligned among them and with
97 representative strains belonging to different NSP4 genotypes according to Matthijnsens et al.
98 (2008a) using Bioedit 7.0.5.3 software (HALL, 1999) and Clustal W 1.83 (THOMPSON et al.,
99 1994) downloaded from the NCBI GenBank database. The strains used were
100 (genotype/accession number/host/strain): a) E1/ GU199490/ Swine/ Gottfried; b) E1/
101 DQ494398/ Bovine/ KJ75; c) E1/ AF144799/ Swine/ A411; d) E1/ D88831/ Swine/ OSU; e)
102 E1/ X69485/ Swine/ YM; f) E1/ U59109/ Human/ M37; g) E2/ AF144805/ Bovine/ B223; h) E3/
103 AF144806/ Canine/ CU 1; i) E4/ AB065285/ Avian/ Ty 1; j) E5/ AF533535/ Lapine/ 160 01;
104 k) E6/ DQ490560/ Human/ RV176 06; l) E7/ U96337/ Murine/ EC; m) E8/ EF442742/ Canine/
105 RV52 96; n) E9/ DQ534017/ Swine/ CMP034; o) E10/ FJ169862/ Avian/ 02V0002G3 and p)
106 E12/ FJ347120/ Bovine/ Arg B383.

107 The nucleotide and amino acid similarities were calculated using Bioedit v. 7.0.5.3
108 software (HALL, 2009). The phylogenetic tree from nucleotide sequences was built using
109 MEGA software version 4 (TAMURA et al., 2007) based on Neighbor-joining method using
110 Maximum Composite Likelihood (1,000 bootstrap trials).

RESULTS

111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135

For the 11 samples, a common fragment of 501 nt (nt 66 to nt 566, using as reference Gottfried Strain accession number GU199490) from RVs NSP4-coding gene was investigated. The strain identification, P and G genotypes, and respective NSP4 gene accession numbers are shown in Table 1. The Genbank accession numbers for the NSP4 partial gene sequences of porcine rotaviruses determined in this study are: HQ840943, HQ840944, HQ840945, HQ840946, HQ840947, HQ840948, HQ840949, HQ840950, HQ840952, HQ840953, HQ840954.

Nucleotide identity ranged from 92.4% to 100% while amino acid ranged from 95.8% to 100%. The comparison of NSP4 genes sequenced in this study with other strains classified as genotype E1 from GenBank revealed a nucleotide identity ranging from 94.4% (strain PORV6 with porcine strains Gottfried and OSU) to 84.1% (strain PORV9 with human strain EF672589) and amino acid identity ranging from 98.8% (strains PORV1; PORV2; PORV3; PORV4; PORV6; PORV7 and PORV11 with Venezuelan porcine strain AF165219) to 89.4% (strain PORV5 with human strain EF672589).

Deduced amino acids of the sequences generated herein revealed a moderate variation among the strains (Fig. 1). Moreover, considering the toxigenic peptide (amino acid 114-135) it was shown that there was a single point mutation on aa 135 presented as alanine in two RV strains and as valine in the other strains. In addition, six other amino acid changes at residues 136 (valine, alanine and serine), 137 (arginine and glycine), 139 (isoleucine and valine), 154 (arginine and lysine), 161 (serine and asparagine) and 174 (serine and proline) were found.

The phylogenetic tree (Fig. 2) depicts that the strains of the present study clustered with E1 genotype representatives, while the others segregated in separate clusters with a resolved genealogy, according to its genotypes.

DISCUSSION

136

137

138 Among non-murine NSP4 amino acid sequences, most of the divergence was observed
139 in the VP4-binding domain (aa 112-148) and in the double-layered particle-binding region (aa
140 161-175) (IOSEF et al., 2002). Strains of the present study showed low degree of polymorphism
141 in both regions, with four and two mutations respectively, as shown in Fig. 1.

142 Strains PORV6 and PORV10 presented amino acid residue alanine at position 135, while
143 the others presented amino acid valine. By comparisons of NSP4 sequences, Zhang et al. (1998)
144 suggested that changes between amino acids 131 and 140 are important for viral pathogenesis,
145 showing that a change from amino acid valine to alanine in the NSP4 protein at this position
146 was important in OSU attenuated strains, as it was associated with loss of the ability of inducing
147 diarrhea in mice, which was also observed in a piglet model with virulent and tissue culture-
148 attenuated human RVs Wa strains (WARD et al., 1996). On the other hand, Kirkwood et al.
149 (1996) found isoleucine at position 135 in symptomatic children, as well as did Mascarenhas et
150 al. (2007).

151 Tyrosine residue at position 131 of NSP4 coding gene has been postulated to be critical
152 for the diarrhegenic activity of the toxic peptide (BALL et al. 1996), but histidine was also
153 found in diarrheic young children (CUNLIFFE et al., 1997; MASCARENHAS et al., 2007).
154 Sequence analysis from porcine strains revealed amino acids serine, alanine and histidine at
155 residue 131 (CIARLET et al., 2000; MATTHIJNSSENS et al., 2010; STEYER et al., 2007). In
156 the present study, all strains showed histidine, as shown in Fig. 1. Therefore, the enterotoxin
157 domain (aa 114-135) is conserved among them, except for one mutation at aa 135.

158 Even though nucleotide and amino acid polymorphism were found both at the toxigenic
159 peptide and VP6-binding domain (aa 112-175) observed in Fig. 1, it was not possible to
160 speculate on the significance of these changes for the virulence of the RV strains since all the

161 animals studied had diarrhea. In other studies, this correlation between virulent and attenuated
162 strains was not observed (ANGEL et al., 1998; WARD et al., 1997), showing the possibility
163 that virus attenuation can occur by several mechanisms, including mutations in other viral
164 proteins. Moreover, the extreme C terminus, including aa methionine at position 175 was shown
165 to be important for double-layered particle (DLP)-binding activity (TAYLOR et al., 1992). As
166 shown in Fig. 1, all the porcine strains presented methionine at this site.

167 This study revealed the occurrence of genotypes G10 and G11 in association with P[6] or
168 P[7] in the swine population. G10 genotype has been widely detected in bovine rotaviruses in
169 Brazil (ALFIERI et al., 2004) and other countries (FALCONE et al., 1999; GARAICOECHEA
170 et al., 2006; HOWE et al., 2008), and also in humans (RAMANI et al., 2009; URASAWA et
171 al., 1993). A study in Thailand also revealed this genotype in pigs (PONGSUWANNA et al.,
172 1996).

173 G11 rotaviruses were first detected in pigs in Mexico and Venezuela (CIARLET et al.,
174 1994; ROSEN et al., 1994; RUIZ et al., 1988) and are believed to be circulating in this
175 population, although in low numbers. In subsequent years, no additional G11 strains were
176 detected in the same or nearby pig farms, but in the last decade, several reports have described
177 the isolation of G11 RVs strains from humans (MATTHIJNSSENS et al., 2010). These authors
178 also showed that multiple reassortment events have occurred between porcine or human G11
179 rotaviruses and co-circulating human Wa-like RVs strains.

180 The phylogenetic tree (Fig. 2) showed that the circulating Brazilian RVs strains belong
181 to E1 genotype, also reported elsewhere in humans, swine, equine, and bovine
182 (MATTHIJNSSENS et al., 2008a), reinforcing the association between E1 genotype and pig
183 RVs previously described. Although evidences for independent segregation of the VP6- and
184 NSP4-encoding genes have been described in porcine RV-A (GHOSH et al., 2006; ITURRIZA-
185 GÓMARA, 2002), considering the limited number of surveyed samples and occurrence of

186 undefined P and G genotypes, it was not possible to observe this pattern among Brazilian
187 samples.

188 Interspecies transmission of rotaviruses may occur in natural and experimental conditions
189 (MARTELLA et al., 2010). The introduction of a new human-animal reassortant RVs strain
190 into the human population could have an impact on the spread of rotavirus disease and also on
191 prevention measures (STEYER et al., 2008). This study also revealed (data not shown) that
192 strain PORV6 had 96,4% amino acid identity with Brazilian strain NB-150, a human strain
193 previously isolated by Mascarenhas et al. (2007) from a newborn with diarrhea who lived in
194 the outskirts of Belém do Pará, Brazil, that reinforce the hypothesis that interspecies
195 transmission may occur naturally, without loss of virulence (VARGHESE et al., 2004).

196 There are numerous examples of RVs interspecies transmission, but there are few
197 documented evidences in which whether the transmission event has involved the whole genome
198 (PALOMBO, 2002). In fact, pigs may serve as a reservoir of RVs for humans, as described by
199 several authors in different countries, such as India, Ecuador and Hungary (BANYAI et al.,
200 2004; BANYAI et al., 2009; VARGHESE et al., 2004). It has been proposed that human RVs
201 Wa-like strains and swine strains have a common origin (MATTHIJNSSENS et al., 2008b),
202 and, recently, a new virus isolated from pigs was closely related to a novel group of human
203 rotaviruses (WAKUDA et al., 2011).

204

205

CONCLUSIONS

206

207 As a conclusion, NSP4 genes of porcine RVs isolated in Brazil during 2008 had only a
208 moderate polymorphism and belonged all to E1, in an extent previously unknown in this
209 country.

210

211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243

ACKNOWLEDGMENTS

This work was supported by grant n° 577884/2008-5 from National Council for Scientific and Technological Development (CNPq) and Brazilian Ministry of Agriculture, Livestock and Food Supply (MAPA).

REFERENCES

ALFIERI, A.F.; ALFIERI, A.A.; BARREIROS, M.A.; LEITE, J.P.; RICHTZENHAIN, L.J. G and P genotypes of group A rotavirus strains circulating in calves in Brazil, 1996-1999. **Veterinary Microbiology**, v.99, n.3-4, p.167-173, 2004.

ANGEL, J.; TANG, B.; FENG, N.; GREENBERG, H.B.; BASS, D. Studies of the role for NSP4 in the pathogenesis of homologous murine rotavirus diarrhea. **Journal of Infectious Diseases**, v.177, n.2, p.455-458, 1998.

BALL, J.M.; TIAN, P.; ZENG, C.Q.; MORRIS, A.P.; ESTES M.K. Age-dependent diarrhea induced by a rotaviral nonstructural glycoprotein. **Science**, v.272, n.5258, p.101-104, 1996.

BANYAI, K.; ESONA, M.D.; KERIN, T.K.; HULL, J.J.; MIJATOVIC, S.; VÁSCONEZ, N.; TORRES, C.; DE FILIPPIS, A.M.; FOYTICH, K.R.; GENTSCH, J.R. Molecular characterization of a rare, human-porcine reassortant rotavirus strain, G11P[6], from Ecuador. **Archives of Virology**, v.154, n.11, p.1823-1829, 2009.

BANYAI, K.; MARTELLA, V.; JAKAB, F.; MELEGH, B.; SZÜCS, G. Sequencing and phylogenetic analysis of human genotype P[6] rotavirus strains detected in Hungary provides evidence for genetic heterogeneity within the P[6] VP4 gene. **Journal of Clinical Microbiology**, v.42, n.9, p.4338-4343, 2004.

CIARLET, M.; HIDALGO, M.; GORZIGLIA, M.; LIPRANDI, F. Characterization of neutralization epitopes on the VP7 surface protein of serotype G11 porcine rotaviruses. **The Journal of General Virology**, v.75, n.8, p.1867-1873, 1994.

CUNLIFFE, N.A.; WOODS, P.A.; LEITE, J.P.G.; DAS, B.K.; RAMACHANDRAN, M.; BHAN, M.K.; HART, C.A.; GLASS, R.I.; GENTSCH, J.R. Sequence analysis of NSP4 gene of human rotavirus allows classification into two main genetic groups. **Journal of Medical Virology**, v.53, n.1, p.41-50, 1997.

FALCONE, E.; TARANTINO, M.; DI TRANI, L.; CORDIOLI, P.; LAVAZZA, A.; TOLLIS, M. Determination of bovine rotavirus G and P serotypes in Italy by PCR. **Journal of Clinical Microbiology**, v.37, n.12, p.3879-3882, 1999.

- 244 GARAICOECHEA, L.; BOK, K.; JONES, L.R.; COMBESSIES, G.; ODEÓN, A.;
245 FERNANDEZ, F.; PARREÑO, V. Molecular characterization of bovine rotavirus circulating
246 in beef and dairy herds in Argentina during a 10-year period (1994-2003). **Veterinary**
247 **Microbiology**, v.118, n.1-2, p.1-11, 2006.
- 248 GHOSH, S.; VARGHESE, V.; SAMAJDAR, S.; BHATTACHARYA, S.K.; KOBAYASHI,
249 N.; NAIK, T.N. Molecular characterization of a porcine group A rotavirus strain with G12
250 genotype specificity. **Archives of Virology**, v.151, n.7, p.1329-1344, 2006.
- 251 GOUVEA, V.; SANTOS, N.; TIMENETSKY, M.C. Identification of bovine and porcine
252 rotavirus G types by PCR. **Journal of Clinical Microbiology**, v.32, n.5, 1338-1340, 1994a.
- 253 GOUVEA, V.; SANTOS, N.; TIMENETSKY, M.C. VP4 typing of bovine and porcine group
254 A rotaviruses by PCR. **Journal of Clinical Microbiology**, v.32, n.5, 1333-1337, 1994b.
- 255 HALL, T.A. BioEdit: a user-friendly biological sequence alignment editor and analysis
256 program for Windows 95/98/NT. **Nucleic Acids Symposium Series**, v.41, n.1, p.95-98, 1999.
- 257 HOWE, L.; SUGIARTO, H.; SQUIRES, R.A. Use of polymerase chain reaction for the
258 differentiation of group A bovine rotavirus G6, G8, and G10 genotypes in the north island of
259 New Zealand. **New Zealand Veterinary Journal**, v.56, n.5, p.218-221, 2008.
- 260 HUANG, H.; SCHROEDER, F.; ESTES, M.K.; MCPHERSON, T.; BALL, J.M. Interaction(s)
261 of rotavirus non-structural protein 4 (NSP4) C- terminal peptides with model membranes. **The**
262 **Biochemical Journal**, v.380, n.3, p.723-733, 2004.
- 263 HYSER, J.M.; UTAMA, B.; CRAWFORD, S.E.; ESTES, M.K. Genetic divergence of
264 rotavirus nonstructural protein 4 results in distinct serogroup-specific viroporin activity and
265 intracellular punctate structure morphologies. **Journal of Virology**, v.86, n.9, p.4921-34, 2012.
- 266 IOSEF, C.; CHANG, K.O.; AZEVEDO, M.S.; SAIF, L.J. Systemic and intestinal antibody
267 responses to NSP4 enterotoxin of Wa human rotavirus in a gnotobiotic pig model of human
268 rotavirus disease. **Journal of Medical Virology**, v.68, n.1, p.119-128, 2002.
- 269 ITURRIZA-GÓMARA, M.; WONG, C.; BLOME, S.; DESSELBERGER, U.; GRAY J.
270 Molecular characterization of VP6 genes of human rotavirus isolates: correlation of genogroups
271 with subgroups and evidence of independent segregation. **Journal of Virology**, v.76, n.13,
272 p.6596-6601, 2002.
- 273 KAPIKIAN, A.Z.; HOSHINO, Y.; CHANOCK, R.M. Rotaviruses. In: KNIPE, D.M.;
274 HOWLEY, P.M.; GRIFFIN, D.E.; LAMB, R.A.; MARTIN, M.A.; ROIZMAN, B.; STRAUS,
275 S.E. (Eds). *Fields Virology*, 4.ed. Philadelphia:Lippincott, Williams & Wilkins, 2001, p.1787-
276 1833.
- 277 KING, A.M.Q.; ADAMS, M.J.; CARSTENS, E.B.; LEFKOWITZ, E.J. Virus taxonomy:
278 classification and nomenclature of viruses: Ninth Report of the International Committee on
279 Taxonomy of Viruses. San Diego: Elsevier Academic Press, 2012, p.603-613.

280 KIRKWOOD, C.D.; COULSON, B.S.; BISHOP, R.F. G3P2 rotaviruses causing diarrheal
281 disease in neonates differ in VP4, VP7 and NSP4 sequence from G3P2 strains causing
282 asymptomatic neonatal infection. *Archives of Virology*, v.141, n.9, p.1661-1676, 1996.

283 LEE, C.N.; WANG, Y.L.; KAO, C.L.; ZAO, C.L.; LEE, C.Y.; CHEN, H.N. NSP4 gene analysis
284 of Rotavirus recovered from infected children with and without diarrhea. **Journal of Clinical
285 Microbiology**, v.38, n.12, p.4471-4477, 2000.

286 MARTELLA, V.; BÁNYAI, K.; MATTHIJNSSENS, J.; BUONAVOGLIA, C.; CIARLET, M.
287 Zoonotic aspects of rotaviruses. **Veterinary Microbiology**, v.140, n.3-4, p.246-255, 2010.

288 MASCARENHAS, J.D.; LEITE, J.P.; LIMA, J.C.; HEINEMANN, M.B.; OLIVEIRA, D.S.;
289 ARAÚJO, I.T.; SOARES, L.S.; GUSMÃO, R.H.; GABBAY, Y.B.; LINHARES, A.C.
290 Detection of a neonatal human rotavirus strain with VP4 and NSP4 genes of porcine origin.
291 **Journal of Medical Microbiology**, v.56, n.4, p.524-532, 2007.

292 MATTHIJNSSENS, J.; CIARLET, M.; HEIMAN, E.; ARIJS, I.; DELBEKE, T.;
293 MCDONALD, S.M.; PALOMBO, E.A.; ITURRIZA-GÓMARA, M.; MAES, P.; PATTON,
294 J.T.; RAHMAN, M.; VAN RANST, M. Full genome-based classification of rotaviruses reveals
295 a common origin between human Wa-Like and porcine rotavirus strains and human DS-1-like
296 and bovine rotavirus strains. **Journal of Virology**, v.82, n.7, p.3204-3219, 2008a.

297 MATTHIJNSSENS, J.; CIARLET, M.; MCDONALD, S.M.; ATTOUI, H.; BÁNYAI, K.;
298 BRISTER, J.R.; BUESA, J.; ESONA, M.D.; ESTES, M.K.; GENTSCH, J.R.; ITURRIZA-
299 GÓMARA, M.; JOHNE, R.; KIRKWOOD, C.D.; MARTELLA, V.; MERTENS, P.P.;
300 NAKAGOMI, O.; PARREÑO, V.; RAHMAN, M.; RUGGERI, F.M.; SAIF, L.J.; SANTOS,
301 N.; STEYER, A.; TANIGUCHI, K.; PATTON, J.T.; DESSELBERGER, U.; VAN RANST, M.
302 Uniformity of rotavirus strain nomenclature proposed by the Rotavirus Classification Working
303 Group (RCWG). **Archives of Virology**, v.156, .8, p.1397-1413, 2011.

304 MATTHIJNSSENS, J.; CIARLET, M.; RAHMAN, M.; ATTOUI, H.; BÁNYAI, K.; ESTES,
305 M. K.; GENTSCH, J. R.; ITURRIZA-GÓMARA, M.; KIRKWOOD, C. D.; MARTELLA, V.;
306 MERTENS, P. P.; NAKAGOMI, O.; PATTON, J. T.; RUGGERI, F. M.; SAIF, L. J.; SANTOS,
307 N.; STEYER, A.; TANIGUCHI, K.; DESSELBERGER, U.; VAN RANST, M.
308 Recommendations for the classification of group A rotaviruses using all 11 genomic RNA
309 segments. Recommendations for the classification of group A rotaviruses using all 11 genomic
310 RNA segments. **Archives of Virology**, v.153, n.8, p.1621-1629, 2008b.

311 MATTHIJNSSENS, J.; OTTO, P.H.; CIARLET, M.; DESSELBERGER, U.; VAN RANST,
312 M.; JOHNE, R. VP6-SEQUENCE-BASED CUTOFF VALUES AS A CRITERION FOR
313 ROTAVIRUS SPECIES DEMARCATION. *ARCHIVES OF VIROLOGY*, V.157, N.6,
314 P.1177-1182, 2012.

315 MATTHIJNSSENS, J.; RAHMAN, M.; CIARLET, M.; ZELLER, M.; HEYLEN, E.;
316 NAKAGOMI, T.; UCHIDA, R.; HASSAN, Z.; AZIM, T.; NAKAGOMI, O.; VAN RANST,
317 M. Reassortment of human rotavirus gene segments into G11 rotavirus strains. **Emerging
318 Infectious Diseases**, v.16, n.4, p.625-630, 2010.

- 320 PALOMBO, E.A. Genetic analysis of Group A rotaviruses, evidence for interspecies
321 transmission of rotavirus genes. **Virus Genes**, v.24, n.1, p.11-20, 2002.
- 322 PONGSUWANNA, Y.; TANIGUCHI, K.; CHIWAKUL, M.; URASAWA, T.; WAKASUGI,
323 F.; JAYAVASU, C.; URASAWA, S. Serological and genomic characterization of porcine
324 rotaviruses in Thailand: detection of a G10 porcine rotavirus. **Journal of Clinical
325 Microbiology**, v.34, n.5, p.1050-1057, 1996.
- 326 RAMANI, S.; ITURRIZA-GOMARA, M.; JANA, A.K.; KURUVILLA, K.A.; GRAY, J.J.;
327 BROWN, D.W.; KANG, G. Whole genome characterization of reassortant G10P[11] strain
328 (N155) from a neonate with symptomatic rotavirus infection: identification of genes of human
329 and animal rotavirus origin. **Journal of Clinical Virology**, v.45, n.3, p.237-244, 2009.
- 330 ROSEN, B.I.; PARWANI, A.V.; LOPEZ, S.; FLORES, J.; SAIF L.J. Serotypic differentiation
331 of rotaviruses in field samples from diarrheic pigs by using nucleic acid probes specific for
332 porcine VP4 and human and porcine VP7 genes. **Journal of Clinical Microbiology**, v.32, n.2,
333 p.311-317, 1994.
- 334 RUIZ, A.M.; LOPEZ, I.V.; LOPEZ, S.; ESPEJO, R.T.; ARIAS, C.F. Molecular and antigenic
335 characterization of porcine rotavirus YM, a possible new rotavirus serotype. **Journal of
336 Virology**, v.62, n.11, p.4331-4336, 1988.
- 337 STEYER, A.; POLJSAK-PRIJATELJ, M.; BARLIC-MAGANJA, D.; JAMNIKAR, U.;
338 MIJOVSKI, J.Z.; MARIN, J. Molecular characterization of a new porcine rotavirus P genotype
339 found in an asymptomatic pig in Slovenia. **Virology**, v.359, n.2, p.275-282, 2007.
- 340 STEYER, A.; POLJSAK-PRIJATELJ, M.; BARLIC-MAGANJA, D.; MARIN, J. Human,
341 porcine and bovine rotaviruses in Slovenia, evidence of interspecies transmission and genome
342 reassortment. **Journal of General Virology**, v.89, n.7, p.1690-1698, 2008.
- 343 TAMURA, K.; DUDLEY, J.; NEI, M.; KUMAR, S. MEGA4: Molecular Evolutionary
344 Genetics Analysis (MEGA) software version 4.0. **Molecular Biology and Evolution**, v.24, n.8,
345 p.1596-1599, 2007.
- 346 TAYLOR, J.A.; MEYER, J.C.; LEGGE, M.A.; O'BRIEN, J.A.; STREET, J.E.; LORD, V.J.;
347 BERGMANN, C.C.; BELLAMY, A.R. Transient expression and mutational analysis of the
348 rotavirus intracellular receptor: the C-terminal methionine residue is essential for ligand
349 binding. **Journal of Virology**, v.66, n.6, p.3566-3572, 1992.
- 350 THOMPSON, J. D.; HIGGINS, D.G.; GIBSON, T.J. CLUSTAL W: improving the sensitivity
351 of progressive multiple sequence alignment through sequence weighting, position-specific gap
352 penalties and weight matrix choice. **Nucleic Acids Research**, v.22, n.22, p.4673-4680, 1994.
- 353 TIAN, P.; ESTES, M.K.; HU, Y.; BALL, J.M.; ZENG, C.Q.; SHILLING, W.P. The rotavirus
354 nonstructural glycoprotein NSP4 mobilizes Ca²⁺ from the endoplasmic reticulum. **Journal of
355 Virology**, v.69, n.9, p.5763-5772, 1995.
- 356

357 TONIETTI, P.O.; HORA, A.S.; SILVA, F.D.F.; RUIZ, V.L.A.; GREGORI, F. Phylogenetic
358 analyses of the VP4 and VP7 genes of porcine group A rotaviruses in Sao Paulo State, Brazil:
359 first identification of G5P[23] in piglets. **Journal of Clinical Microbiology**, v.51, n.8, p.2750-
360 2753, 2013.

361 URASAWA, T.; TANIGUCHI, K.; KOBAYASHI, N.; MISE, K.; HASEGAWA, A.;
362 YAMAZI, Y.; URASAWA, S. Nucleotide sequence of VP4 and VP7 genes of a unique human
363 rotavirus strain Mc35 with subgroup I and serotype 10 specificity. **Virology**, v.195, n.2, p.766-
364 771, 1993.

365 VARGHESE, V.; DAS, S.; SINGH, N.B.; KOJIMA, K.; BHATTACHARYA, S.K.;
366 KRISHNAN, T.; KOBAYASHI, N.; NAIK, T.N. Molecular characterization of a human
367 rotavirus reveals porcine characteristics in most of the genes including VP6 and NSP4.
368 **Archives of Virology**, v.149, n.1, p.155-172, 2004.

369 WAKUDA, M.; IDE, T.; SASAKI, J.; KOMOTO, S.; ISHII, J.; SANEKATA, T.;
370 TANIGUCHI, K. Porcine rotavirus closely related to novel group of human rotaviruses.
371 **Emerging Infectious Diseases**, v.17, n.8, p.1491-1493, 2011.

372 WARD, R.L.; MASON, B.B.; BERNSTEIN, D.I.; SANDER, D.S.; SMITH, V.E.; ZANDLE,
373 G.A.; RAPPAPORT, R.S. Attenuation of a human rotavirus vaccine candidate did not correlate
374 with mutations in the NSP4 protein gene. **Journal of Virology**, v.71, n.8, p.6267-6270, 1997.

375 WARD, L.A.; ROSEN, B.I.; YUAN, L.; SAIF, L.J. Pathogenesis of an attenuated and a virulent
376 strain of group A human rotavirus in neonatal gnotobiotic pigs. **The Journal of General**
377 **Virology**, v.77, n.7, 1431-1441, 1996.

378 ZHANG, M.; ZENG, C.Q.; DONG, Y.; BALL, J.M.; SAIF, L.J.; MORRIS, A.P.; ESTES, M.K.
379 mutations in rotavirus nonstructural glycoprotein NSP4 are associated with altered virus
380 virulence. **Journal of Virology**, v.72, n.5 p.3666-3672, 1998.

381

382

TABLE 1

383

Strain	Genotype P	Genotype G	NSP4 Accession number
PORV1	P[6]	G[11]	HQ840943
PORV2	P[6]	G[11]	HQ840944
PORV3	P[6]	G[11]	HQ840945
PORV4	P[6]	G[11]	HQ840946
PORV5	-	-	HQ840947
PORV6	-	G[10]	HQ840948
PORV7	P[7]	-	HQ840949
PORV8	P[6]	-	HQ840952
PORV9	P[7]	-	HQ840950
PORV10	P[7]	G[10]	HQ840953
PORV11	-	G[10]	HQ840954

384


385 Table 1: P and G genotypes and accession numbers of partial NSP4 sequences RVs from
 386 piglets samples in São Paulo State, Brazil. Gaps indicate genotypes that were not defined.

387

388

FIGURE 1

389



390 Fig. 1. Section of the alignment of the deduced 175 amino acids (aa 9-175) of the NSP4-coding
391 gene from rotavirus detected in porcine stool samples from Brazilian herds. The marked area
392 refers to the toxigenic peptide (residues 114 to 135).


393

394

395

FIGURE 2

396



397

398 Fig. 2. Unrooted neighbor-joining tree for a stretch of 501 nucleotides (nt 25-525) of the NSP4-
399 coding gene, showing the proposed E genotypes. Taxa designated as "PORV (1 to 11)" are
400 related to the Brazilian field strains from the present study; numbers at each node are the
401 bootstrap values greater than 50% obtained with 1,000 replicates.